



Radio sources

P. Charlot

Laboratoire d'Astrophysique de Bordeaux





Outline



- **Introduction**
 - Continuum and spectral line emission processes
 - The radio sky: galactic and extragalactic
- **History of radioastronomy**
 - The first 50 years (1932-1981)
- **Active Galactic Nuclei (AGN)**
 - Observational properties
 - Standard unified model
- **Imaging radiosources**
 - Aperture synthesis
 - The case of VLBI

Electromagnetic emission can be divided into two types:


- **Continuum emission**

- emission over a very broad frequency range
- usually due to the acceleration of charged particles moving with a wide-range of energy



- **Spectral line emission**



- emission over a very narrow frequency range
- usually due to the discrete transitions in the internal energy states of atoms or molecules






- Thermal emission


- Black body radiation for objects with temperature $T \sim 3-30$ K
- Bremsstrahlung (free-free) emission: deflection of a charged particle (electron) in the electric field of another charged particle (ion)






- Non-thermal emission

- Synchrotron radiation: relativistic electrons spiraling around weak magnetic field lines







- Neutral hydrogen (21 cm)
 - spin-flip transition between high-energy state and low-energy state of the H atom (aligned vs opposed spins for p+ and e-)
- Molecular lines (CO, CS, CN,...)
 - Produced by changes in the vibrational or rotational states of their electrons (due to collisions or interactions)
- Maser emission
 - Amplification of incident radiation passing through clouds of gas




Ionized gas in the Orion nebula




Betelgeuse




Masers around the star TX Cam




SS 433 (X-ray binary)



Pulsars







Supernova remnant





Credits: M. Kramer (pulsar animation) - all other images courtesy of NRAO/AUI

Atomic hydrogen emission



Continuum emission




Images courtesy of NRAO/AUI





History of radioastronomy

When it all started...

- 1932: Karl Jansky discovers cosmic radio waves while investigating sources of radio noise adversely affecting transatlantic communications




Images courtesy of NRAO/AUI




Galactic centre


1944: Grote Reber builds the first parabolic radio telescope and makes the first map of the radio sky (160 MHz & 480 MHz)




Images courtesy of NRAO/AUI



- Later on detects radio emission from Cas-A, Cyg-A, Cyg-X,...
- Multi-frequency observations reveal non-thermal emission





- 1949: identification of two strong radio sources (Cen-A, Virgo A) with nearby galaxies (Bolton et al.)
- 1954: identification of the radio source Cyg-A with a distant galaxy (Baade & Minkowski)



→ reveals the extragalactic nature of some of the radio sources

- 1963: discovery of quasars (quasi-stellar radio source)

- Identification of 3C273 with a faint 13th magnitude star-like source
- ... but with emission lines shifted to longer wavelengths by 16%




Schmidt (1963)


- ➔ most distant known object in the Universe at the time but also intrinsically the most luminous one.
- ➔ first member of a new class of objects now referred to as « Active Galactic Nuclei » (AGN)

- 1967: discovery of pulsars

- Periodic source of radio emission with $T=2s$
- associated with dense fast rotating neutron star



Credit: M. Kramer



Bell & Hewish (1968)

P. Charlot

- Apparent faster-than-light motions in AGN (known as superluminal motions)
 - 1971: through visibility curves (Whitney et al.)
 - 1981: through VLBI imaging (Pearson et al.)

- Interpreted as a geometrical effect in a relativistically-expanding source





Image courtesy of NRAO/AUI




Active Galactic Nuclei (AGN)

- AGN are located at cosmological distances
- Distance is measured by redshift: $z = \lambda - \lambda_0 / \lambda_0$





Credit: Gene Smith



Schneider et al. (2007)

- highest-redshift quasar known at present is at $z=7.085$ while the highest-redshift radio source is at $z=6.21$
- AGN have no detected proper motions


Relative Variation of Thermal and Non-thermal Radiation Emissions




- Non-thermal emission
- About 15-20% of AGNs are « radio-loud » while the rest are « radio-quiet »

AGN may vary at any wavelength (from radio to gamma-rays) on timescales of minutes to years


Years




Months



Years




Hours



Aharonian et al. (2008)


Intrinsic fast variations imply very small physical size for the variable region

- An object that shows variability on a timescale Δt cannot be larger than $c \Delta t$.



Credit: Gene Smith

- Variability on a scale of a few minutes means that the AGN size cannot be larger than a few light-minutes.



 Point-like


 One sided

 Two-sided


A sample of X band (8 GHz) VLBI maps with milliarcsecond resolution picked up randomly from the *Bordeaux VLBI Image Database (BVID)*

3C120


2000



2001




2003




Credit:
Bordeaux VLBI
Image Database

2234+282


2000




2001




2003




2007







Credit: Craig Walker



Credit: MOJAVE database




Credit: MOJAVE database



Credit: MOJAVE database

Major components

- Black hole
- Accretion disk
- Torus
- Pair of relativistic jets




http://heasarc.gsfc.nasa.gov/docs/objects/agn/agn_model.html

Credit: C. M. Urry & P. Padovani

Object with jet close to the plane of the sky

- weak core
- two-sided jet



Polatidis et al. (1999)

Object with jet pointing towards the observer

- strong core
- one-sided jet






Image courtesy of NRAO/AUI and R. C. Walker

- Dichotomy radio-loud/
radio quiet
- Classification according
to viewing angle
 - Radio loud: BL Lac, quasars,
radio galaxies
 - Radio-quiet: QSO, Seyfert 1,
Seyfert 2




Credit: C. M. Urry & P. Padovani




Credit: Alan Marscher


2 GHz




8 GHz



24 GHz





43 GHz



Credit: Radio Reference Frame Image Database

➔ Source structure becomes more compact at higher frequencies



SI = 1
Excellent

SI = 4
very bad

Source structure effect in VLBI delay

The **structure index** (SI) – defined as the median « source structure effect » over the u-v plane – indicates the astrometric suitability of the sources.



Imaging radiosources

Complex visibility


$$V(u,v) = A \exp(i\Phi) = \iint_s I(x,y) \exp(-2\pi i(ux+vy)) dx dy$$

- u,v (measured in wavelengths) are spatial frequencies in E-W and N-S directions, i.e. the baseline length
- x,y (measured in radians) are angles in tangent plane relative to a reference position in the E-W and N-S directions

Sky brightness

$$I(x,y) = \iint_{\text{plan}(u,v)} V(u,v) \exp(2\pi i(ux+vy)) du dv$$


- The complex visibility is the 2D Fourier transform of the brightness on the sky




Credit: David Wilner

- Amplitude tells « how much » of a certain frequency component
- Phase tells where this component is located





$I(x,y)$



$A\{V(u,v)\}$




$\Phi\{V(u,v)\}$




Credit: David Wilner

- Amplitude tells « how much » of a certain frequency component
- Phase tells where this component is located





$I(x,y)$



$A\{V(u,v)\}$



$\Phi\{V(u,v)\}$



Credit: David Wilner



Aperture synthesis: ideal case




In practice, the (u,v) plane is incompletely covered and only a limited number of spatial frequencies is sampled

$$I(x,y) = \sum_{j=1,N} V(u_j, v_j) \exp(2\pi i(u_j x + v_j y))$$

→ Image distortion

→ Needs deconvolution




Credit: Tim Cornwell



Aperture synthesis: example




Source model




Credit: Tim Cornwell

u-v plane



Amplitudes







Aperture synthesis: example




u-v plane



Dirty beam



Dirty beam (2-D)




Credit: Tim Cornwell




Aperture synthesis: example




Source model



Dirty map





CLEAN map



Credit: Tim Cornwell



Another case

Model



“Dirty” beam

“Dirty” image




CLEAN image

Credit: Tim Cornwell

- Φ cannot be used directly
- Assumes $\Phi_s = 0$ to start (point source)
- Takes advantage of closure phases:

$$\Phi_{1-2} + \Phi_{2-3} + \Phi_{3-1} = f(I(x,y))$$
- Iterates to adjust Φ_s





VLBI imaging surveys



- **Bordeaux VLBI Image Database (2 & 8 GHz)**
<http://www.obs.u-bordeaux1.fr/BVID/>
- **Radio Reference Frame Image Database (2 & 8 GHz)**
<http://www.usno.navy.mil/USNO/astrometry/vlbi-products/rrfid>
- **VLBA Calibrator Survey (2 & 8 GHz)**
<http://www.vlba.nrao.edu/astro/calib/index.shtml>
- **MOJAVE data base (15 GHz)**
<http://www.physics.purdue.edu/astro/MOJAVE/index.html>
- **VLBI Imaging and Polarimetry Survey (5 & 15 GHz)**
<http://www.phys.unm.edu/~gbtaylor/VIPS/vipscat/vipsncapindx.shtml>



Acknowledgements



Information and figures presented in this lecture have been taken from the following sources:

- Mike Garrett's radioastronomy course
http://www.astron.nl/~mag/dokuwiki/doku.php?id=radio_astronomy_course_description
- NRAO Synthesis Imaging Workshops (2002-2012)
<http://www.aoc.nrao.edu/events/synthesis/2012/>
- NRAO image gallery
<http://images.nrao.edu/>
- Bordeaux VLBI Image Database
<http://www.obs.u-bordeaux1.fr/BVID/>
- Radio Reference Frame Image Database
<http://www.usno.navy.mil/USNO/astrometry/vlbi-products/rfid>
- MOJAVE data base
<http://www.physics.purdue.edu/astro/MOJAVE/index.html>